

SCIENTIFIC REPORTS



OPEN

Very rapid long-distance sea crossing by a migratory bird

José A. Alves^{1,2}, Maria P. Dias^{3,4}, Verónica Méndez⁵, Borgný Katrínardóttir⁶ & Tómas G. Gunnarsson²

Received: 06 September 2016

Accepted: 04 November 2016

Published: 30 November 2016

Landbirds undertaking within-continent migrations have the possibility to stop *en route*, but most long-distance migrants must also undertake large non-stop sea crossings, the length of which can vary greatly. For shorebirds migrating from Iceland to West Africa, the shortest route would involve one of the longest continuous sea crossings while alternative, mostly overland, routes are available. Using geolocators to track the migration of Icelandic whimbrels (*Numenius phaeopus*), we show that they can complete a round-trip of 11,000 km making two non-stop sea crossings and flying at speeds of up to 24 m s^{-1} ; one of the fastest recorded for shorebirds flying over the ocean. Although wind support could reduce flight energetic costs, whimbrels faced headwinds up to twice their ground speed, indicating that unfavourable and potentially fatal weather conditions are not uncommon. Such apparently high risk migrations might be more common than previously thought, with potential fitness gains outweighing the costs.

Recent advances in tracking movements of individual birds are revolutionising our understanding of avian migration^{1,2}. New tracking technologies (e.g. geolocators and PTT transmitters) have revealed migratory journeys in excess of 5000 km of active flight^{1–3}, setting the endurance exercise record of any animal⁴. Ultra long-distance continuous migratory flights have been suggested to facilitate avoidance of predators, parasites and pathogens⁵, but such continuous exercise is also known to increase mortality risk⁴. Migratory landbirds typically avoid crossing large ecological barriers such as mountain-ranges, deserts and oceans^{6,7}, often using routes over suitable habitats along which stopping to rest and refuel is possible, for example by following the edge of continental land-masses⁶. However, when detours from the shortest route considerably increase travel distance, energy or time, crossing ecological barriers may be beneficial¹. Birds breeding at high latitudes are often those that undertake the longest non-stop flights, crossing oceans^{1,8–10}, ice-caps¹¹ and deserts², but such long flights can incur high mortality risk due to exhaustion associated with prolonged unfavourable weather conditions⁵. Species migrating by flying long distances continuously over oceanic waters^{1,3,8,12} and land masses² all undertake stop-overs during either both journeys^{2,3,8,12} or pre-nuptial migration only^{1,6}. During pre-nuptial migration, the use of stop-over sites might allow gauging conditions closer to the Arctic and subarctic breeding grounds as arriving too early can also be disadvantageous if conditions are unsuitable upon arrival⁶. Additionally, by refuelling during migration, birds can restore body reserves thus increasing the likelihood of reaching the breeding grounds in good condition and at the right time^{6,13}. Non-stop long distance flights over unsuitable habitats entail considerable survival risk as no sheltering options are available and future weather conditions in distant locations are impossible to predict¹⁴. Indeed the largest mortality events recorded on migration (5000–200 000 individuals^{15–18}) refer to landbirds crossing large waterbodies and encountering adverse weather conditions⁶. However, favourable winds can also play a fundamental role in the crossing over large expanses of unsuitable habitat such as oceans⁶. Many species wait for tailwinds at coastal sites for several days before embarking on large sea crossings, departing only when wind subsidies are considerable^{6,14}. If significant headwinds are encountered during oceanic flights, birds must endure substantially longer flight periods which can result in depletion of fat and muscles and exhaustion¹⁹. It is therefore expected that birds depart for long distance sea crossings with favourable tailwinds and that flight air speeds are as high as possible, independent of winds encountered *en route*, in order to reduce exposure to potentially adverse conditions.

¹DBIO & CESAM-Centre for Environmental and Marine Studies, University of Aveiro, Aveiro, Portugal. ²South Iceland Research Centre, University of Iceland, Fjölheimar IS-800 Selfoss & IS-861 Gunnarsholt, Iceland. ³BirdLife International, The David Attenborough Building, Pembroke Street, Cambridge, CB2 3QZ, UK. ⁴MARE-Marine and Environmental Sciences Centre, ISPA-Instituto Universitário, Rua Jardim do Tabaco 34, 1149-041, Lisboa, Portugal. ⁵School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK. ⁶Ecology Department, Icelandic Institute of Natural History, 210, Gardabaer, Iceland. Correspondence and requests for materials should be addressed to J.A.A. (email: j.alves@uea.ac.uk)



Figure 1. Icelandic whimbrel (*Numenius phaeopus islandicus*) carrying a geolocator (left: photo by Tómas G. Gunnarsson) attached to a leg flag (right: photo by Camilo Carneiro).




Figure 2. Geotracked migratory routes of four Icelandic whimbrels (*Numenius phaeopus islandicus*) between breeding sites in Iceland and wintering areas in West Africa during post-nuptial/autumn (left) and pre-nuptial/spring (right) migration. Track sections are coloured as a function of ground speed. All individuals flew non-stop in Autumn whilst two made a stop-over in the UK or Ireland in Spring (details in Table 1). Maps created using R 3.1.2 using packages ggplot2, ggmap, raster and RgoogleMaps³³ (image data providers: US Dept. of State Geographer © 2016 Google) in WSG 48 coordinate reference system.

Iceland hosts important populations of several migratory wader (or shorebird) species²⁰ which winter in Europe and west Africa^{21,22}. These species have to negotiate one of the longest continuous sea crossing of all Arctic and subarctic breeding landbirds⁶. Species migrating to west Africa (e.g. the Bijagós archipelago in Guinea-Bissau) could either undertake a non-stop oceanic flight (~5800 km) or, after an initial sea crossing (~800 km) to the UK, follow the continental land masses to the wintering grounds (~5200 km). Both alternatives result in similar distances and some Icelandic breeding waders are known to follow the coastline²¹, even those for which a single flight overwater is potentially feasible²³. The extent to which either of these routes is used is not known and although the non-stop oceanic flight is potentially of higher risk, this will likely depend on wind conditions during migration. We deployed geolocator tags on Icelandic whimbrels (*Numenius phaeopus islandicus* Brehm, 1831), a species which is known to winter in West Africa²² to investigate (1) if non-stop flights over oceanic waters between Iceland and West Africa are undertaken during autumn and spring migration; (2) the level of wind support encountered *en route* and how this affects flight speed.

Results and Discussion

Ten adult breeding Icelandic whimbrels were tagged with geolocators in June 2012 (Fig. 1), seven were recorded on their territories in June 2013, of which four tags with data were retrieved. During the post-nuptial migration in autumn 2012, all four whimbrels flew non-stop to their wintering areas in west Africa (Fig. 2), covering distances of ~3900 to 5500 km in 5 days (Table 1) and, on occasion, achieving some of the fastest recorded speeds for terrestrial birds on long-distance flight over oceanic waters (up to 18–24 m s⁻¹). During the return migration, two of these birds stopped for 11 (Male 2) and 15 days (Female 2), covering a total distance of ~10,500 and 11,000 km, respectively. However, the remaining female and male completed the return migration in another

Migration	Autumn				Spring			
	Male 1	Male 2	Female 1	Female 2	Male 1	Male 2	Female 1	Female 2
Onset of migration (departure)	03-Aug	06-Aug	06-Aug	03-Aug	20-Apr	22-Apr	29-Apr	23-Apr
End of migration (arrival)	07-Aug	10-Aug	10-Aug	07-Aug	25-Apr	10-May	04-May	14-May
Total duration (d)	5	5	5	5	6	19	6	22
Stopover time (d)	0	0	0	0	0	11	0	15
Total migration distance (km)	5425	5171	3898	5535	5555	5364	3865	5560
Total duration (h)	107.8	107.9	78.9	120.0	121.2	444.1	127.3	540.6
Total migration speed (km h ⁻¹)	50.31	47.94	49.38	46.14	45.83	12.08	30.37	10.29
Non-stop flights (over ground speed)								
Max speed (m s ⁻¹)	24.18	18.60	17.93	21.91	19.71	21.29	13.45	18.19
Min speed (m s ⁻¹)	5.75	9.89	5.66	5.36	3.57	2.19	3.08	4.79
Average (m s ⁻¹)	15.55	14.38	13.32	14.35	13.87	8.24	9.09	9.63
Sd	6.84	3.00	4.19	6.42	5.10	5.40	3.17	4.25
N	7	8	8	9	10	14	11	15

Table 1. Timings, distances and speed of total migration and non-stop flights by Icelandic whimbrels that flew direct to the winter grounds (all birds in Autumn) and that made a stop-over (Male 2 and Female 2) or a direct flight (Male 1 and Female 1) in the return migration (Spring).

continuous flight (Fig. 2) for a total round trip of ~7800 (Female 1) and 11,000 (Male 1) km, respectively (Table 1). Completing such a long annual migration cycle in two long-distance flights is highly unusual for species where such studies have been undertaken. Only one other long-distance Arctic migrant wader, the Pacific Golden Plover (*Pluvialis fulva*), has been recorded undertaking such non-stop flights (ca. 9700 km total distance) between Alaska and Hawaii⁹. For this species however, no alternative route over coastal land masses is possible without considerably increasing flight distance.

In order to assess how wind conditions encountered on these journeys varied, particularly for flights with and without stopovers, we quantified wind support at each position just prior to and during the migratory flight. All four whimbrels departed Iceland in favourable wind conditions (i.e. tailwinds), but all four arrived in West Africa having faced headwinds, mostly in the later part of the journey (Fig. 3). Conversely, only one individual departed from the winter grounds in favourable wind conditions (Male 1, the earliest to depart), while all others departed in headwinds of 1.3–4.8 m s⁻¹, with one individual encountering headwinds for virtually the entire journey, including after stopping-over (Female 2, Fig. 3). A pre-nuptial migration strategy involving a stopover is likely safer regarding potentially unfavourable weather conditions encountered *en route* and upon arrival, but stopping to refuel will reduce overall migration speed. Indeed, and despite all tracked individuals departing the winter grounds at similar times (22nd–29th of April), those flying non-stop arrived before those that undertook a stopover. This includes the last individual to depart which flew non-stop (Female 1) and arrived at the breeding grounds 6 to 10 days before the two individuals that made a stopover (of 11–15 days), thus overtaking its conspecifics²³. The earlier arrival of the female and male that flew non-stop to Iceland could be advantageous if they capitalize by nesting early, as this is known to increase breeding success, particularly in Arctic and subarctic systems²⁴. However, laying dates did not differ substantially between the individuals undertaking direct flights (20th–31st of May) or those that made a stopover (25th May–7th of June), suggesting that timing of breeding is constrained by other or additional factors, such as environmental conditions for nesting or timing of mate arrival (Supplementary Fig. S1).

The very fast ground speeds achieved by migrating whimbrels were influenced by the wind speeds encountered *en route*, particularly at altitudes of 1500 m (Table 2). Wind support at this altitude accounts for 4 to 36% of fastest speeds of each individual, with the highest wind assistance corresponding to the maximum recorded ground speed of 24.2 m s⁻¹ (87 km h⁻¹). Some individuals also reach very fast speeds whilst facing headwinds which can be 2 to 40% of their ground speed, and in its most extreme case resulting in airspeed of 25.0 m s⁻¹ (90 km h⁻¹; Female 2). Average speeds for the entire continuous migratory flight are similar to those of other species crossing oceans (50–65 km h⁻¹^{5,8,9}) which are also strongly influenced by wind speed *en route*⁵. By flying at high speeds and non-stop over open ocean these species reduce the time on migration and might be using an “airspace corridor” to avoid predators, parasites and pathogens⁵. But if wind conditions at distant locations along the route are unfavourable, such migratory strategy can result in mortality by exhaustion, even after arriving at destination (TGG, pers. observation).

Our ability to track migration is providing new insights into the extraordinary capacity of birds to move extremely fast over very large distances, by continuously sustaining endurance exercise during several days. Wind support is crucial during such extreme journeys, but current predictions of changes in climatic patterns, specifically changes on regional scale wind patterns²⁵, can potentially have a considerably disproportionate negative effect on those species that regularly undertake non-stop long distance flights over unsuitable habitats. Variation in migratory strategies within the same population will likely allow coping with potential changes, but predicting such responses requires an understanding on how these migration strategies can arise and are maintained. In addition, by linking different migratory strategies to associated fitness consequences will be key in our ability to anticipate demographic changes for migratory populations.




Figure 3. Variation in wind speed (m s^{-1}) experienced during autumn (left column) and spring (right column) migratory flights of four Icelandic whimbrels. Each row is one individual, following Table 1, from top to bottom: Male 1, Female 1, Male 2, Female 2. Negative wind values (blue) indicate headwind and positive values indicate tailwind. Wind was estimated for three flight altitudes: 1500, 3000 and 5500 m. Distance from departure location was converted into percentage for ease of interpretation.

	Estimate (SE)	df	t value	p value
Intercept	12.11 (0.56)	71	21.487	<0.001
Wind 1500 m	0.51 (0.21)	71	2.436	0.017
Wind 3000 m	0.16 (0.29)	71	-1.197	0.235
Wind 5500 m	0.70 (0.13)	71	1.433	0.156

Table 2. Results of GLMM of wind speed at three altitudes on ground speed of four Icelandic whimbrels tracked during migration between Iceland and West Africa.

Methods

Bird tracking. Given Icelandic whimbrels previously established adult return rate (ca. 60–80%)²⁶, in June 2012 we deployed 10 geolocators (Intigeo W65A9R), Migrate Technology) on breeding birds in South Iceland ($63^{\circ} 47' \text{N}$, $20^{\circ} 12' \text{W}$). All individuals were caught with a nest trap (Moudry TR60; www.moudry.cz), ringed with metal and colour rings and measured to determine sex²⁶. Seven of these birds were recorded breeding in the same location 12 months later, five of which (two females and three males) were re-captured and the geolocator collected. All of these individuals were from different pairs and the geolocator of one male was corrupted as a result of saltwater entering the device, leaving four individuals available for analysis. All animal handling and protocols were carried out in accordance with relevant guidelines and regulations under licenses issued by Icelandic (Natural History Institute; license number 365) and International regulatory bodies (International Wader Study Group; license number 1235).

Positional data and flight speed. Light data from the geolocators were smoothed twice²⁷ and used to estimate positions²⁸ during migration (i.e. between Iceland and West Africa) using IntiProc (v. 1.03, Migrate

Technology, Ltd.) and “GeoLight” package in R²⁹, assuming a sun elevation angle of -6° based on *in situ* geolocator calibration prior to deployment. Total migration length, distance (great circle route) and speed were estimated between the last and first positions on land in the breeding areas (Iceland) or wintering areas (W Africa). As geolocator positions are only attained at a minimum of 12 hours intervals, flight speed (time taken to cover the distance between two sequential positions-in m s^{-1}) was estimated for each 12 hour flight segment defined as two sequential positions between the first location outside the breeding, wintering or stop-over areas and the first location on land (at breeding, wintering or stop-over areas).

Wind support. Wind data at the location (± 2.5 degrees) and time of each geolocator recorded position was extracted from NOAA using the dataset Reanalysis by NCEP³⁰. Headwind or tailwind vector between sequential positions was interpolated for three altitudes (1500, 3000 and 5500 m) using function “NCEP.interp” from package RNCEP³¹. To test at which altitude wind speed had an effect on ground speed, we built a GLMM with individual as random factor to control the non-independence and DFs were calculated using the Satterthwaite approximation in package lmerTest. All analysis and calculations were performed in R 2.15.0³².

References

- Battley, P. F. *et al.* Contrasting extreme long-distance migration patterns in bar-tailed godwits *Limosa lapponica*. *J. Avian Biol.* **43**, 21–32 (2012).
- Klaassen, R. H. G. *et al.* Great flights by great snipes: long and fast non-stop migration over benign habitats. *Biol. Lett.* **7**, 833–5 (2011).
- Niles, L. J. *et al.* First results using light level geolocators to track Red Knots in the Western Hemisphere show rapid and long intercontinental flights and new details of migration pathways. *Wader Study Gr. Bull.* **117**, 123–130 (2010).
- Piersma, T. Why marathon migrants get away with high metabolic ceilings: towards an ecology of physiological restraint. *J. Exp. Biol.* **214**, 295–302 (2010).
- Gill, R. E. *et al.* Extreme endurance flights by landbirds crossing the Pacific Ocean: ecological corridor rather than barrier? *Proc. Biol. Sci.* **276**, 447–57 (2009).
- Newton, I. *The Migration Ecology of Birds*. (Academic Press, 2008).
- Alerstam, T. *Bird Migration*. (Cambridge University Press, 1990).
- Minton, C. D. T. *et al.* Initial results from light level geolocator trials on Ruddy Turnstone *Arenaria interpres* reveal unexpected migration route. *Wader Study Gr. Bull.* **117**, 9–14. (2010).
- Johnson, O. W. *et al.* Tracking the migrations of Pacific Golden-Plovers (*Pluvialis fulva*) between Hawaii and Alaska: New insight on flight performance, breeding ground destinations, and nesting from birds carrying light level geolocators. *Wader Study Gr. Bull.* **118**, 26–31 (2011).
- Bairlein, F. *et al.* Cross-hemisphere migration of a 25 g songbird. *Biol. Lett.* **8**, 505–507 (2012).
- Fox, A. D., Glahder, C. M. & Walsh, A. J. Spring migration routes and timing of Greenland white-fronted geese—results from satellite telemetry. *Oikos* **103**, 415–425 (2003).
- Minton, C. *et al.* New insights from geolocators deployed on waders in Australia. *Wader Study Gr. Bull.* **120**, 37–46 (2013).
- Alerstam, T. & Lindström, Å. In *Bird Migr. Physiol. Ecophysiol.* (ed. Gwinner, E.) 331–351 (Springer Berlin Heidelberg), doi: 10.1007/978-3-642-74542-3_22 (1990).
- Shamoun-Baranes, J. *et al.* Stochastic atmospheric assistance and the use of emergency staging sites by migrants. *Proc. Biol. Sci.* **277**, 1505–11 (2010).
- James, P. Destruction of warblers on Padre Island, Texas. *Wilson Bull.* **68**, 224–227 (1951).
- Webster, F. S. The spring migration, April 1–May 31, 1974, South Texas region. *Am. Birds* **28**, 822–825 (1974).
- Jansson, R. B. The spring migration, April 1–May 31, 1976, Western Great Lakes region. *Am. Birds* **30**, 844–846 (1976).
- Alerstam, T. Findings of dead birds drifted ashore reveal catastrophic mortality among early spring migrants, especially Rooks *Corvus frugilegus*, over the southern Baltic sea. *Anser* **27**, 181–218 (1988).
- Pennycuik, C. J. & Battley, P. F. Burning the engine: a time-marching computation of fat and protein consumption in a 5420-km non-stop flight by great knots, *Calidris tenuirostris*. *Oikos* **103**, 323–332 (2003).
- Gunnarsson, T. G. *et al.* Large-scale habitat associations of birds in lowland Iceland: Implications for conservation. *Biol. Conserv.* **128**, 265–275 (2006).
- Thorisson, B., Eyjólfsson, V., Gardarsson, A., Albertsdóttir, H. B. & Gunnarsson, T. G. The non-breeding distribution of Icelandic Common Ringed Plovers. *Wader Study Gr. Bulletin* **119**, 97–101 (2012).
- Gunnarsson, T. G. & Guðmundsson, G. a. Migration and non-breeding distribution of Icelandic Whimbrels *Numenius phaeopus islandicus* as revealed by ringing recoveries. *Wader Study* **123**, 44–48 (2016).
- Alves, J. A. *et al.* Overtaking on migration: does longer distance migration always incur a penalty? *Oikos* **121**, 464–470 (2012).
- Gunnarsson, T. G., Gill, J. A., Newton, J., Potts, P. M. & Sutherland, W. J. Seasonal matching of habitat quality and fitness in a migratory bird. *Proc. Biol. Sci.* **272**, 2319–23 (2005).
- IPCC. Intergovernmental Panel on Climate Change: fifth Assessment Report. (2014).
- Katrínardóttir, B., Pálsson, S., Gunnarsson, T. G. & Sigurjónsdóttir, H. Sexing Icelandic Whimbrels *Numenius phaeopus islandicus* with DNA and biometrics. *Ringed Migr.* **28**, 43–46 (2013).
- Pütz, K. Spatial and temporal variability in the foraging areas of breeding king penguins. *Condor* **104**, 528–538 (2002).
- Dias, M. P., Granadeiro, J. P., Phillips, R. A., Alonso, H. & Catry, P. Breaking the routine: individual Cory’s shearwaters shift winter destinations between hemispheres and across ocean basins. *Proc. Biol. Sci.* **278**, 1786–93 (2011).
- Lisovski, S. & Hahn, S. GeoLight-processing and analysing light-based geolocator data in R. *Methods Ecol. Evol.* **3**, 1055–1059 (2012).
- Kalnay, E. *et al.* The NCEP/NCAR 40-Year Reanalysis Project. *Bull. Am. Meteorol. Soc.* **77**, 437–471 (1996).
- Kemp, M. U., Emiel van Loon, E., Shamoun-Baranes, J. & Bouten, W. RNCEP: global weather and climate data at your fingertips. *Methods Ecol. Evol.* **3**, 65–70 (2012).
- R Development Core Team, R. R: A Language and Environment for Statistical Computing. *R Found. Stat. Comput.* **1**, 409 (2011).
- Loecher, M. & Ropkins, K. RgoogleMaps and loa : Unleashing R Graphics Power on Map Tiles. *J. Stat. Softw.* **63**, 1–18 (2015).

Acknowledgements

For support with fieldwork in Iceland and discussions, Graham Appleton, Jennifer Gill, Juan Carlos Illera and Sara Pardal. This work was funded by NERC, RANNIS - The Icelandic Research Fund (Grant of Excellence 130412-051) and FCT (individual grant to JAA: SFRH/BPD/91527/2012). NCEP Reanalysis data provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from their Web site at <http://www.esrl.noaa.gov/psd/>.

Author Contributions

J.A.A. conceived the study with input from T.G.G. and M.P.D., M.P.D. sourced the geolocators and B.K. and T.G.G. attained permissions. J.A.A., B.K., T.G.G. and V.M. developed fieldwork and J.A.A. & M.P.D. analysed the data. J.A.A. lead the writing with substantial contributions from M.P.D., B.K., V.M. and T.G.G.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Alves, J. A. *et al.* Very rapid long-distance sea crossing by a migratory bird. *Sci. Rep.* **6**, 38154; doi: 10.1038/srep38154 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016

SCIENTIFIC REPORTS

OPEN

Corrigendum: Very rapid long-distance sea crossing by a migratory bird

José A. Alves, Maria P. Dias, Verónica Méndez, Borgný Katrínardóttir & Tómas G. Gunnarsson

Scientific Reports 6:38154; doi: 10.1038/srep38154; published online 30 November 2016; updated on 28 April 2017

The original version of this Article contained an error. The authors reported the maximum flight speed of Whimbrels (*Numenius phaeopus*) as the fastest recorded for shorebirds flying over the ocean, but this was not the case.

As a result, in the Abstract,

“Using geolocators to track the migration of Icelandic whimbrels (*Numenius phaeopus*), we show that they can complete a round-trip of 11,000 km making two non-stop sea crossings and flying at speeds of up to 24 m s^{-1} ; the fastest recorded for shorebirds flying over the ocean”.

now reads:

“Using geolocators to track the migration of Icelandic whimbrels (*Numenius phaeopus*), we show that they can complete a round-trip of 11,000 km making two non-stop sea crossings and flying at speeds of up to 24 m s^{-1} ; one of the fastest recorded for shorebirds flying over the ocean”.

In the Results and Discussion section,

“During the post-nuptial migration in autumn 2012, all four whimbrels flew non-stop to their wintering areas in west Africa (Fig. 2), covering distances of ~3900 to 5500 km in 5 days (Table 1) and, on occasion, achieving the fastest recorded speeds for terrestrial birds on long-distance flight over oceanic waters (up to $18\text{--}24 \text{ m s}^{-1}$)”.

now reads:

“During the post-nuptial migration in autumn 2012, all four whimbrels flew non-stop to their wintering areas in west Africa (Fig. 2), covering distances of ~3900 to 5500 km in 5 days (Table 1) and, on occasion, achieving some of the fastest recorded speeds for terrestrial birds on long-distance flight over oceanic waters (up to $18\text{--}24 \text{ m s}^{-1}$)”.

These errors have now been corrected in the PDF and HTML versions of the Article.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017