

Real-World Six- and Twelve-Month Drug Retention, Remission, and Response Rates of Secukinumab in 2,017 Patients With Psoriatic Arthritis in Thirteen European Countries

Brigitte Michelsen,¹  Stylianos Georgiadis,² Daniela Di Giuseppe,³ Anne G. Loft,⁴ Michael J. Nissen,⁵ Florenzo Iannone,⁶  Manuel Pombo-Suarez,⁷ Herman Mann,⁸ Ziga Rotar,⁹ Kari K. Eklund,¹⁰ Tore K. Kvien,¹¹  Maria J. Santos,¹²  Bjorn Gudbjornsson,¹³ Catalin Codreanu,¹⁴ Sema Yilmaz,¹⁵ Johan K. Wallman,¹⁶  Cecilie H. Brahe,¹⁷ Burkhard Möller,¹⁸  Ennio G. Favalli,¹⁹ Carlos Sánchez-Piedra,²⁰ Lucie Nekvindova,²¹ Matija Tomsic,⁹ Nina Trokovic,¹⁰ Eirik K. Kristianslund,¹¹ Helena Santos,²² Thorvardur J. Löve,²³ Ruxandra Ionescu,¹⁴ Yavuz Pehlivan,²⁴ Gareth T. Jones,²⁵  Irene van der Horst-Bruinsma,²⁶ Lykke M. Ørnbjerg,¹⁷ Mikkel Østergaard,²⁷ and Merete L. Hetland²⁸

Objective. There is a lack of real-life studies on interleukin-17 (IL-17) inhibition in psoriatic arthritis (PsA). We assessed real-life 6- and 12-month effectiveness (i.e., retention, remission, low disease activity [LDA], and response rates) of the IL-17 inhibitor secukinumab in PsA patients overall and across 1) number of prior biologic/targeted synthetic disease-modifying antirheumatic drugs (b/tsDMARDs), 2) years since diagnosis, and 3) European registries.

Methods. Thirteen quality registries in rheumatology participating in the European Spondyloarthritis Research Collaboration Network provided longitudinal, observational data collected as part of routine care for secondary use. Data were pooled and analyzed with Kaplan-Meier plots, log rank tests, Cox regression, and multiple linear and logistic regression analyses.

Results. A total of 2,017 PsA patients started treatment with secukinumab between 2015 and 2018. Overall secukinumab retention rates were 86% and 76% after 6 and 12 months, respectively. Crude (LUNDEX adjusted) 6-month remission/LDA (LDA including remission) rates for the 28-joint Disease Activity Index for Psoriatic Arthritis, the Disease Activity Score in 28 joints using the C-reactive protein level, and the Simplified Disease Activity Index (SDAI) were 13%/46% (11%/39%), 36%/55% (30%/46%), and 13%/56% (11%/47%), and 12-month rates were 11%/46% (7%/31%), 39%/56% (26%/38%), and 16%/62% (10%/41%), respectively. Clinical Disease Activity Index remission/LDA rates were similar to the SDAI rates. Six-month American College of Rheumatology 20%/50%/70% improvement criteria responses were 34%/19%/11% (29%/16%/9%); 12-month rates were 37%/21%/11% (24%/14%/7%). Secukinumab effectiveness was significantly better for b/tsDMARD-naïve patients, similar across time since diagnosis (<2/2–4/>4 years), and varied significantly across the European registries.

Conclusion. In this large real-world study on secukinumab treatment in PsA, 6- and 12-month effectiveness was comparable to that in previous observational studies of tumor necrosis factor inhibitors. Retention, remission, LDA, and response rates were significantly better for b/tsDMARD-naïve patients, were independent of time since diagnosis, and varied significantly across the European countries.

INTRODUCTION

Psoriatic arthritis (PsA) is a heterogeneous inflammatory rheumatic disease affecting, e.g., peripheral joints, axial spine, skin, and entheses, with significant impact on health-related quality of life

(1–3). The treatment options for PsA have improved during the last few decades with the introduction of biologic disease-modifying antirheumatic drugs (bDMARDs) and targeted synthetic DMARDs (tsDMARDs) (4). Nevertheless, a recent real-world study of >14,000 patients with PsA, who started treatment with a tumor

The EuroSpA collaboration was financially supported by Novartis. Novartis had no influence on the data collection, statistical analyses, manuscript preparation, or decision to submit.

¹Brigitte Michelsen, MD, Cand. Mag., PhD: Rigshospitalet, Glostrup, Denmark, Hospital of Southern Norway Trust, Kristiansand, Norway, and

Diakonhjemmet Hospital, Oslo, Norway; ²Stylianos Georgiadis, PhD: Rigshospitalet, Glostrup, Denmark; ³Daniela Di Giuseppe, PhD: Karolinska Institutet, Stockholm, Sweden; ⁴Anne G. Loft, MD, PhD: DANBIO Registry and Rigshospitalet, Glostrup, Denmark, and Aarhus University Hospital, Aarhus, Denmark; ⁵Michael J. Nissen, MD, PhD: Geneva University Hospital, Geneva,

SIGNIFICANCE & INNOVATIONS

- Secukinumab retention, remission, low disease activity (LDA), and response rates were significantly better for biologics-naïve patients after 6 as well as 12 months of treatment.
- Overall 6- and 12-month secukinumab retention rates were high; remission, LDA, and response rates were good; and overall effectiveness was comparable to that in previous observational studies of tumor necrosis factor inhibitors.
- This study is to date the largest real-world study on secukinumab effectiveness in patients with psoriatic arthritis, including 2,017 patients from 13 European national registries.
- The study documents the effectiveness of secukinumab for treatment of psoriatic arthritis in clinical practice and shows significantly better outcomes for biologics-naïve patients. This may be taken into consideration in treatment decisions in routine clinical care.

necrosis factor inhibitor (TNFi), showed that less than one-half of the patients had achieved clinical remission after 6 months (5). Thus, there is an unmet need for other treatment options in patients with PsA (2,6).

The fully human IgG monoclonal interleukin-17A (IL-17A) inhibitor secukinumab was approved for use in PsA patients in the European Union in 2015 (7). Secukinumab has demonstrated good efficacy and safety in randomized controlled trials (RCTs) (8–10), whereas large observational studies on its effectiveness in patients with PsA are lacking.

Hence, the main objective of this study was to assess the overall real-life 12-month retention rate of secukinumab in PsA patients in Europe. Secondary objectives were to assess the overall 6-month secukinumab retention rate and 6- and 12-month remission, low disease activity (LDA), and response rates. These aims were assessed overall, as well as compared across number of previous b/tsDMARD treatments, time since diagnosis, and the European registries.

Switzerland; ⁶Florenzo Iannone, MD, PhD: GISEA Registry and University of Bari, Bari, Italy; ⁷Manuel Pombo-Suarez, MD, PhD: Hospital Clínico Universitario, Santiago de Compostela, Spain; ⁸Herman Mann, MD, PhD: Charles University, Prague, Czech Republic; ⁹Ziga Rotar, MD, PhD, Matija Tomsic, MD, PhD: University Medical Centre Ljubljana, Ljubljana, Slovenia; ¹⁰Kari K. Eklund, MD, PhD, Nina Trokovic, MD, PhD: Helsinki University Hospital and ORTON Orthopaedic Hospital of the Orton Foundation, Helsinki, Finland; ¹¹Tore K. Kvien, MD, PhD, Eirik K. Kristianslund, MD, PhD: Diakonhjemmet Hospital, Oslo, Norway; ¹²Maria J. Santos, MD, PhD: Reuma.pt Registry and Universidade de Lisboa, Lisbon, Portugal; ¹³Bjorn Gudbjornsson, MD, PhD: Centre for Rheumatology Research (ICEBIO), University Hospital and University of Iceland, Reykjavik, Iceland; ¹⁴Catalin Codreanu, MD, PhD, Ruxandra Ionescu, MD, PhD: University of Medicine and Pharmacy Carol Davila, Bucharest, Romania; ¹⁵Sema Yilmaz, MD, PhD: Selcuk University School of Medicine, Selcuklu, Turkey; ¹⁶Johan K. Wallman, MD, PhD: Lund University, Skåne University Hospital, Lund, Sweden; ¹⁷Cecilie H. Brahe, MD, PhD, Lykke M. Ørnberg, MD, PhD: Rigshospitalet and DANBIO Registry, Glostrup, Denmark; ¹⁸Burkhard Möller, MD, PhD: Universitätsklinik für Rheumatologie, Immunologie und Allergologie, Inselspital, Bern, Switzerland; ¹⁹Ennio G. Favalli, MD, PhD: ASST Gaetano Pini-CTO Institute, Milan, Italy; ²⁰Carlos Sánchez-Piedra, MD, PhD: Spanish Society of Rheumatology, Madrid, Spain; ²¹Lucie Nekvindova, MD, PhD: Charles University, Prague, Czech Republic, and Institute of Biostatistics and Analyses, Masaryk University, Brno, Czech Republic; ²²Helena Santos, MD, PhD: Reuma.pt Registry and Portuguese Institute of Rheumatology, Lisbon, Portugal; ²³Thorvardur J. Löve, MD, PhD: University of Iceland and Landspítali University Hospital, Reykjavik, Iceland; ²⁴Yavuz Pehlivan, MD, PhD: Uludağ University, Bursa, Turkey; ²⁵Gareth T. Jones, PhD: University of Aberdeen, Aberdeen, UK; ²⁶Irene van der Horst-Bruinsma, MD, PhD: Amsterdam University Medical Centres, VU University Medical Centre, Amsterdam, The Netherlands; ²⁷Mikkel Østergaard, MD, PhD, DMSc: Rigshospitalet, Glostrup, Denmark, and University of Copenhagen, Copenhagen, Denmark; ²⁸Merete L. Hetland, MD, PhD, DMSc: Rigshospitalet and DANBIO Registry, Glostrup, Denmark, and University of Copenhagen, Copenhagen, Denmark.

Drs. Østergaard and Hetland contributed equally to this work.

Dr. Michelsen has received consulting fees from Novartis (less than \$10,000) and research support from Novartis. Dr. Georgiadis has received research support from Novartis. Dr. Loft has received consulting fees and/or speaking fees from AbbVie, Eli Lilly and Company, Janssen, MSD, Novartis, Pfizer, and UCB (less than \$10,000 each). Dr. Nissen has received consulting fees and/or speaking fees from AbbVie, Celgene, Eli Lilly and Company, Novartis, and Pfizer (less than \$10,000 each). Dr. Iannone has received consulting fees and/or speaking fees from AbbVie, Eli Lilly and Company, MSD, Novartis,

Pfizer, Roche, Janssen, Bristol Myers Squibb, and Sanofi (less than \$10,000 each). Dr. Rotar has received consulting fees and/or speaking fees from AbbVie, Amgen, Biogen, Eli Lilly and Company, Medis, MSD, Novartis, Pfizer, and Sanofi (less than \$10,000 each). Dr. Kvien has received consulting fees and/or speaking fees from AbbVie, Amgen, Biogen, Celltrion, Egis, Eli Lilly and Company, Eva Pharma, Ewopharma, Gilead, Hikma, MSD, Mylan, Novartis/Sandoz, Oktal, Orion Pharma, Hospira/Pfizer, Roche, Sanofi, and UCB (less than \$10,000 each) and research support to Diakonhjemmet Hospital from AbbVie, Bristol Myers Squibb, MSD, Pfizer, Roche, and UCB. Dr. Santos has received speaking fees from AbbVie, Novartis, Pfizer, and Roche (less than \$10,000 each). Dr. Gudbjornsson has received speaking fees from Novartis (less than \$10,000). Dr. Codreanu has received consulting fees and/or speaking fees from AbbVie, Accord Healthcare, Alfasigma, Egis, Eli Lilly and Company, Ewopharma, Genesis, Mylan, Novartis, Pfizer, Roche, Sandoz, and UCB (less than \$10,000 each). Dr. Wallman has received consulting fees from Celgene, Eli Lilly and Company, and Novartis (less than \$10,000 each). Dr. Brahe has received research support from Novartis. Dr. Favalli has received consulting fees and/or speaking fees from AbbVie, Bristol Myers Squibb, Eli Lilly and Company, UCB Novartis, Pfizer, and Sanofi-Genzyme (less than \$10,000 each). Dr. Ionescu has received consulting fees and/or speaking fees from AbbVie, Amgen, Eli Lilly and Company, Ewopharma, Novartis/Sandoz, Pfizer, Roche, and UCB (less than \$10,000 each). Dr. Jones has received research support from AbbVie, Pfizer, UCB, Amgen, and GlaxoSmithKline. Dr. van der Horst-Bruinsma has received consulting fees and/or speaking fees from AbbVie, Eli Lilly and Company, Novartis, UCB, Bristol Myers Squibb, MSD, UCB, and Pfizer (less than \$10,000 each). Dr. Ørnberg has received research support from Novartis. Dr. Østergaard has received consulting fees and/or speaking fees from AbbVie, Bristol Myers Squibb, Boehringer-Ingelheim, Celgene, Eli Lilly and Company, Hospira, Janssen, Merck, Novartis, Novo, Orion, Pfizer, Regeneron, Roche, Sandoz, Sanofi, and UCB (less than \$10,000 each) and research support from AbbVie, Bristol Myers Squibb, Celgene, Merck, and Novartis. Dr. Hetland has received consulting fees and/or speaking fees from Eli Lilly and Company, Orion Pharma, Biogen, Pfizer, CellTrion, Merck, and Samsung Bioepis (less than \$10,000 each) and research support from Bristol Myers Squibb, MSD, AbbVie, Roche, Novartis, Biogen, and Pfizer. No other disclosures relevant to this article were reported.

Address correspondence to Brigitte Michelsen, MD, Cand. Mag., PhD, Copenhagen Center for Arthritis Research, Center for Rheumatology and Spine Diseases, Centre for Head and Orthopaedics, Rigshospitalet, Valdemar Hansens Vej 17, 2600 Glostrup, Denmark. Email: brigitte.michelsen@regionh.dk.

Submitted for publication August 28, 2020; accepted in revised form January 12, 2021.

PATIENTS AND METHODS

The European Spondyloarthritis Research Collaboration Network (EuroSpA RCN). The EuroSpA RCN currently includes 15 European quality registries of spondyloarthritis patients (5,11,12). The collaboration was initiated in 2016, but data collection had started as early as 1999 in some of the registries. The main aim of the collaboration is to investigate clinically relevant research questions by secondary use of prospectively collected real-life data (5,11,12). All data are anonymized in the different registries before upload to a secured central server. The data are quality checked and pooled prior to statistical analyses.

Patients. The studies in the EuroSpA collaboration are based on secondary use of real-world data already collected in the different registries, i.e., independently of the current study. In this study, we included data from PsA patients starting secukinumab for the first time between May 2015 and December 2018 in 13 countries in the EuroSpA RCN (ranked by number of patients): ARTIS (Sweden), DANBIO (Denmark), SCQM (Switzerland), GISEA (Italy), BIOBADASER (Spain), ATTRA (Czech Republic), biorx.si (Slovenia), Reuma.pt (Portugal), NOR-DMARD (Norway), ROB-FIN (Finland), ICEBIO (Iceland), RRBR (Romania), and TURKBIO (Turkey). Inclusion criteria for the current analyses were age ≥ 18 years at treatment initiation, a diagnosis of PsA as judged by the treating rheumatologist, and a registered start and, if relevant, stop date of secukinumab. The exclusion criterion was patients with no available clinical data.

Assessments. We included data on age, sex, time since diagnosis, current smoking status (yes/no), body mass index (kg/m^2), start and stop dates of secukinumab, previous b/tsDMARD treatment, evaluator's global assessment, patient's global assessment, pain and fatigue, C-reactive protein (CRP) level (mg/liter), erythrocyte sedimentation rate (ESR, mm/hour), 28-joint Disease Activity Index for Psoriatic Arthritis (DAPSA28) score (13), Disease Activity Score in 28 joints using the CRP level (DAS28-CRP) score (14), Clinical Disease Activity Index (CDAI) score (15), and Simplified Disease Activity Index (SDAI) score (15). The following remission/LDA and response measures were calculated at 6 and 12 months treatment: DAPSA28 remission (≤ 4) (13), DAPSA28 LDA (≤ 14) (13), DAS28-CRP remission (< 2.6) (16), DAS28-CRP LDA (≤ 3.2) (17), CDAI remission (≤ 2.8) (15), CDAI LDA (≤ 10) (15), SDAI remission (≤ 3.3) (15), SDAI LDA (≤ 11) (15), American College of Rheumatology (ACR)/European Alliance of Associations for Rheumatology (EULAR) Boolean remission (18), change in DAPSA28, DAS28-CRP, CDAI, and SDAI, ACR 20%/50%/70% improvement criteria (ACR20/50/70) response (19), and EULAR response (moderate/good) (17).

Primary and secondary outcomes. Primary outcome was the overall 12-month secukinumab retention rate. Secondary

outcomes were the overall 6-month secukinumab retention rate and 6- and 12-month remission, LDA, and response rates.

Statistical analyses. All statistical analyses were performed according to a predefined statistical analysis plan developed by the researchers in the EuroSpA collaboration. Descriptive statistics were performed for demographic data and baseline disease activity measures. All effectiveness analyses were compared across 1) the number of previous b/tsDMARDs ($0/1/\geq 2$), 2) years since diagnosis ($< 2/2-4/> 4$), and 3) the individual registries. Drug retention was explored by Kaplan-Meier analyses with log rank test and by Cox regression analyses adjusted for age, sex, and time since diagnosis (comparisons 1 and 3 above), or age and sex (comparison 2 above).

Remission, LDA, response rates, and change measures were compared by chi-square test, Fisher's exact test, and Kruskal-Wallis test, as appropriate, as well as by multiple linear and logistic regression analyses adjusted for age, sex, and time since diagnosis (comparisons 1 and 3 above), or age and sex (comparison 2 above), as appropriate. Multiple comparisons for the number of previous b/tsDMARDs ($0/1/\geq 2$) were performed by log rank test, chi-square test, Fisher's exact test, and Kruskal-Wallis with post hoc Dunn test, as appropriate, where *P* values were adjusted by applying the Holm's correction.

Significance of relevant groups was tested through likelihood ratio test or Wald test, as appropriate, by comparing 2 nested models. A significance level of 0.05 was used for all statistical tests. In adjusted analyses, multivariate imputation by chained equations (including 50 imputed data sets) was used for 463 patients with missing data for time since diagnosis (no missing data for age and sex). The variables used for imputing time since diagnosis were age, sex, country, and b/tsDMARD treatment series number. None of the other variables including outcome was imputed. To avoid inflating remission and response rates, these were provided both as crude values and with LUND-DEX (20) adjustment, i.e., integrating clinical response and adherence to therapy in a composite value. In the Kaplan-Meier and Cox regression analyses, observations were censored by first occurrence of 1 of the following: end of registry follow-up or date of data extraction. Patients who stopped treatment due to remission or other reasons (e.g., pregnancy) were censored at the stop date to reflect that their withdrawal was not due to lack of effectiveness or adverse events. The baseline date was defined as the secukinumab treatment start date. To assess the robustness regarding the main outcomes, sensitivity analyses for patients 1) having ≥ 1 swollen joints (of 28) at baseline and 2) having date of data extraction at least 12 months after secukinumab treatment start were performed. Competing risk analysis was performed for a cumulative incidence curve showing withdrawal due to adverse events and lack of effectiveness. Numbers available for each of the analyses are shown in Supplementary Tables 1–7,

available on the *Arthritis Care & Research* website at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560>. Statistical analyses were performed with R, version 3.6.1.

Ethics. Approval of the study was obtained from the respective national data protection agencies and research ethical committees according to the individual legal regulatory requirements in the different registries/countries. The study was performed in accordance with the Declaration of Helsinki and followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines (21).

RESULTS

We included a total of 2,017 PsA patients who started secukinumab for the first time (Table 1). The number of patients included from the different European registries varied from 30 (TURKBIO) to 657 (ARTIS). Significant heterogeneity in demographic data and baseline disease activity across the European

registries was found (see Supplementary Table 1, available at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560>). Information on doses was not registered systematically. Of 745 patients in whom doses were registered, 42% of the patients initiated secukinumab 150 mg, and 58% initiated secukinumab 300 mg.

Secukinumab retention rates. The crude 95% confidence interval secukinumab retention rates were overall 76% (74–78%) after 12 months and 86% (85–88%) after 6 months of treatment (Table 2). Secukinumab retention rates after 6 as well as 12 months of treatment were significantly higher in biologics-naïve patients compared with patients previously treated with ≥ 2 b/tsDMARDs (Table 2 and Figure 1A). The findings were similar in 6- and 12-month adjusted Cox regression analyses (see Supplementary Table 8, available on the *Arthritis Care & Research* website at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560>).

Secukinumab retention was not significantly associated with time since diagnosis, either in unadjusted or in adjusted analyses

Table 1. Demographic characteristics and baseline disease activity measures*

	All patients (n = 2,017)	b/tsDMARD naïve (n = 441)	1 prior b/tsDMARD (n = 461)	≥ 2 prior b/tsDMARDs (n = 1,115)	P†
Age, years	52 (44–60)	50 (41–58)	51 (44–59)	53 (45–60)	<0.001
Men, %	43	51	46	39	<0.001
Years since diagnosis	7 (3–13)	4 (1–10)	6 (2–12)	8 (5–14)	<0.001
Current smokers, %	19	18	22	18	0.356
BMI, kg/m ²	27.5 (24.3–31.2)	28.1 (24.1–31.8)	27.3 (24.1–30.1)	27.3 (24.5–31.6)	0.309
B/tsDMARD treatment, % first (% last previous)					<0.001 (<0.001)
Adalimumab	29 (21)	–	30 (30)	28 (18)	
Certolizumab	5 (8)	–	5 (5)	5 (10)	
Etanercept	28 (22)	–	25 (25)	29 (20)	
Golimumab	10 (12)	–	9 (9)	10 (13)	
Infliximab	22 (13)	–	15 (15)	25 (12)	
Other‡	7 (24)	–	15 (15)	3 (27)	
CRP, mg/liter	5 (2–12)	7 (2–19)	4 (2–9)	5 (2–12)	<0.001
ESR, mm/hour	16 (7–31)	20 (8–36)	13 (6–27)	16 (7–30)	0.002
TJC28	4 (1–9)	5 (1–10)	3 (1–8)	4 (1–9)	<0.001
SJC28	1 (0–4)	2 (0–6)	1 (0–3)	2 (0–4)	<0.001
Patient global score	70 (50–83)	70 (51–84)	67 (42–80)	70 (50–85)	<0.001
Pain score	66 (46–80)	65 (45–78)	62 (40–78)	68 (48–81)	<0.001
Fatigue score	70 (50–85)	65 (50–80)	65 (41–80)	73 (55–87)	<0.001
Evaluator global score	40 (20–60)	57 (30–75)	35 (20–50)	35 (20–50)	<0.001
HAQ score	1.1 (0.6–1.6)	1.0 (0.5–1.5)	1.0 (0.5–1.4)	1.2 (0.8–1.8)	<0.001
DAPSA28 score	25.9 (17.4–37.6)	29.1 (19.1–41.9)	22.3 (13.5–32.4)	26.2 (18.0–37.6)	<0.001
DAS28-CRP score	4.2 (3.2–5.0)	4.5 (3.6–5.4)	3.8 (2.7–4.6)	4.2 (3.3–5.0)	<0.001
SDAI score	19.5 (12.9–28.9)	24.4 (15.3–35.4)	16.9 (10.0–24.3)	18.9 (13.0–27.5)	<0.001
CDAI score	18.0 (12.0–26.7)	22.6 (14.3–33.9)	16.0 (8.9–23.6)	17.5 (12.0–25.4)	<0.001

* Values are the median (interquartile range) unless indicated otherwise. Numbers available for each of the analyses are shown in Supplementary Table 5, available on the *Arthritis Care & Research* website at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560>. b/tsDMARD = biologic/targeted synthetic disease-modifying antirheumatic drug; BMI = body mass index; CDAI = Clinical Disease Activity Index; CRP = C-reactive protein; DAPSA28 = 28-joint Disease Activity Index for Psoriatic Arthritis; DAS28-CRP = Disease Activity Score in 28 joints using the CRP level; ESR = erythrocyte sedimentation rate; HAQ = Health Assessment Questionnaire; SDAI = Simplified Disease Activity Index; SJC28 = Swollen joint count in 28 joints; TJC28 = Tender joint count in 28 joints.

† Comparisons between b/tsDMARD-naïve patients and 1 prior and ≥ 2 prior b/tsDMARD-treated patients were performed with Kruskal-Wallis or chi-square test, as appropriate.

‡ Other previous b/tsDMARDs include ustekinumab, rituximab, abatacept, tocilizumab, apremilast, anakinra, and additionally (never as first b/tsDMARD) baricitinib and tofacitinib. Patients were included between May 2015 and December 2018.

Table 2. Treatment effectiveness after 6 and 12 months of secukinumab treatment (unadjusted analyses)*

	All patients (n = 2,017)	b/tsDMARD naive (n = 441)	1 prior b/tsDMARD (n = 461)	≥2 prior b/tsDMARDs (n = 1,115)	P†
Secukinumab drug retention rate, % (95% CI)					
6 months	86 (85–88)	90 (87–93)	86 (83–90)	85 (83–87)	0.045§
12 months	76 (74–78)	82 (78–86)	78 (74–82)	72 (70–75)	0.001§
Time in weeks to secukinumab withdrawal before 12 months due to the following‡					
Primary and secondary lack of effectiveness	24 (17, 33)	24 (17, 35)	24 (17, 30)	24 (17, 34)	0.691
Adverse events	14 (6, 28)	22 (13, 28)	15 (7, 25)	12 (5, 29)	0.395
Remission	21 (20, 43)	20 (19, 20)	–	43 (32, 43)	0.236
Other reasons	21 (12, 32)	27 (15, 40)	10 (4, 36)	21 (15, 27)	0.161
DAPSA28 score					
6 months	15.1 (8.2, 25.0)	10.1 (5.2, 17.5)	15.7 (9.0, 22.0)	16.9 (9.6, 27.1)	<0.001¶
12 months	14.9 (8.1, 24.8)	10.2 (4.1, 16.3)	15.2 (8.4, 23.6)	16.3 (10.0, 26.0)	<0.001¶
DAS28-CRP score					
6 months	3.0 (2.2, 4.0)	2.5 (1.9, 3.3)	3.1 (2.2, 3.9)	3.2 (2.4, 4.2)	<0.001#
12 months	3.0 (2.2, 4.0)	2.5 (1.7, 3.3)	3.0 (2.1, 3.9)	3.2 (2.4, 4.2)	<0.001¶
SDAI score					
6 months	10.2 (5.4, 16.7)	6.9 (3.5, 11.0)	10.4 (6.3, 15.3)	11.4 (6.6, 18.5)	<0.001¶
12 months	9.2 (5.2, 15.2)	5.7 (2.5, 9.5)	9.3 (5.8, 16.2)	10.5 (6.8, 16)	<0.001¶
CDAI score					
6 months	9.3 (4.9, 15.9)	6.2 (3.4, 10.5)	9.4 (5.5, 14.4)	10.9 (6.0, 17.8)	<0.001#
12 months	8.5 (4.4, 14.2)	5.1 (2.1, 9.3)	8.7 (5.2, 14.6)	9.8 (5.8, 14.9)	<0.001¶
Change in DAPSA28 score from baseline					
6 months	–9.5 (–20.7, –0.2)	–17.2 (–27.5, –8.3)	–8.5 (–17.6, –0.1)	–6.6 (–18.3, 0.3)	<0.001¶
12 months	–10.3 (–21.9, –1.3)	–16.2 (–28.0, –8.3)	–5.0 (–10.6, 1.0)	–10.3 (–21.9, –0.2)	<0.001#
Change in DAS28-CRP score from baseline					
6 months	–0.9 (–1.9, –0.1)	–2.0 (–3.0, –1.1)	–0.8 (–1.7, 0.1)	–0.6 (–1.6, 0.01)	<0.001¶
12 months	–1.1 (–2.0, –0.1)	–1.9 (–3.1, –1.0)	–0.5 (–1.3, 0.03)	–1.0 (–1.9, –0.02)	<0.001#
Change in SDAI score from baseline					
6 months	–8.9 (–17.4, –2.0)	–16.9 (–26.1, –9.3)	–7.5 (–13.5, –1.1)	–6.0 (–13.4, –0.2)	<0.001¶
12 months	–9.7 (–18.6, –2.4)	–15.0 (–24.2, –7.5)	–4.9 (–10.4, 1.3)	–9.6 (–17.9, –2.2)	<0.001#
Change in CDAI score from baseline					
6 months	–8.0 (–16.1, –1.6)	–15.1 (–24.6, –8.0)	–6.0 (–13.1, –1.4)	–5.3 (–12.2, –0.1)	<0.001¶
12 months	–8.8 (–16.0, –2.0)	–13.9 (–21.5, –7.3)	–5.0 (–10.4, 0.8)	–8.1 (–15.9, –1.5)	<0.001#
DAPSA28 score ≤4, %					
6 months					
Crude	13	23	13	10	<0.001§
LUNDEX adjusted‡	11	20	11	8	<0.001¶
12 months					
Crude	11	22	11	8	<0.001¶
LUNDEX adjusted‡	7	17	7	5	<0.001§
DAPSA28 score ≤14, %					
6 months					
Crude	46	64	45	41	<0.001¶
LUNDEX adjusted‡	39	57	37	34	<0.001¶
12 months					
Crude	46	70	46	40	<0.001¶
LUNDEX adjusted‡	31	52	30	26	<0.001¶
DAS28-CRP score <2.6, %					
6 months					
Crude	36	53	35	30	<0.001¶
LUNDEX adjusted‡	30	47	29	25	<0.001¶
12 months					
Crude	39	55	41	34	<0.001¶
LUNDEX adjusted‡	26	41	27	21	<0.001¶
DAS28-CRP score ≤3.2, %					
6 months					
Crude	55	71	57	49	<0.001¶
LUNDEX adjusted‡	46	63	47	40	<0.001¶
12 months					
Crude	56	72	55	51	<0.001¶
LUNDEX adjusted‡	38	54	37	33	<0.001¶

(Continued)

Table 2. (Cont'd)

	All patients (n = 2,017)	b/tsDMARD naive (n = 441)	1 prior b/tsDMARD (n = 461)	≥2 prior b/tsDMARDs (n = 1,115)	P†
SDAI score ≤3.3, %					
6 months					
Crude	13	24	13	9	<0.001¶
LUNDEX adjusted‡	11	21	11	8	<0.001¶
12 months					
Crude	16	32	11	11	<0.001¶
LUNDEX adjusted‡	10	24	8	7	<0.001¶
SDAI score ≤11, %					
6 months					
Crude	56	75	56	48	<0.001¶
LUNDEX adjusted‡	47	66	47	39	<0.001¶
12 months					
Crude	62	81	58	56	<0.001¶
LUNDEX adjusted‡	41	61	39	36	<0.001¶
CDAI score ≤2.8, %					
6 months					
Crude	13	19	12	10	0.004§
LUNDEX adjusted‡	10	17	10	8	0.002§
12 months					
Crude	16	32	14	11	<0.001¶
LUNDEX adjusted‡	11	24	10	7	<0.001¶
CDAI score ≤10, %					
6 months					
Crude	55	74	58	46	<0.001¶
LUNDEX adjusted‡	46	66	48	38	<0.001¶
12 months					
Crude	59	79	58	53	<0.001¶
LUNDEX adjusted‡	40	59	39	34	<0.001¶
ACR/EULAR Boolean remission, %					
6 months					
Crude	9	20	8	6	<0.001¶
LUNDEX adjusted‡	8	18	6	5	<0.001¶
12 months					
Crude	9	17	9	6	<0.001§
LUNDEX adjusted‡	6	12	6	4	<0.001§
ACR20 response, %					
6 months					
Crude	34	59	26	27	<0.001¶
LUNDEX adjusted‡	29	52	22	22	<0.001¶
12 months					
Crude	37	63	16	33	<0.001¶
LUNDEX adjusted‡	24	47	10	21	<0.001¶
ACR50 response, %					
6 months					
Crude	19	41	11	13	<0.001¶
LUNDEX adjusted‡	16	36	9	11	<0.001¶
12 months					
Crude	21	45	4	16	<0.001¶
LUNDEX adjusted‡	14	34	3	10	<0.001¶
ACR70 response, %					
6 months					
Crude	11	26	7	6	<0.001¶
LUNDEX adjusted‡	9	23	6	5	<0.001¶
12 months					
Crude	11	28	4	6	<0.001¶
LUNDEX adjusted‡	7	21	3	4	<0.001¶
EULAR good/moderate response, %					
6 months					
Crude	59	83	57	50	<0.001¶
LUNDEX adjusted‡	49	74	48	41	<0.001¶

(Continued)

Table 2. (Cont'd)

	All patients (n = 2,017)	b/tsDMARD naive (n = 441)	1 prior b/tsDMARD (n = 461)	≥2 prior b/tsDMARDs (n = 1,115)	P†
12 months					
Crude	60	79	44	59	<0.001¶
LUNDEX adjusted‡	40	59	30	38	<0.001¶

* Values are the median (interquartile range) unless indicated otherwise. Numbers available for each of the analyses are shown in Supplementary Table 6, available on the *Arthritis Care & Research* website at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560>. 95% CI = 95% confidence interval; ACR = American College of Rheumatology; ACR20/50/70 = ACR 20%/50%/70% improvement criteria; b/tsDMARD = biologic/targeted synthetic disease-modifying antirheumatic drug; CDAI = Clinical Disease Activity Index; CRP = C-reactive protein; DAPSA28 = 28-joint Disease Activity Index for Psoriatic Arthritis; DAS28-CRP = Disease Activity Score in 28 joints using the CRP level; EULAR = European Alliance of Associations for Rheumatology; SDAI = Simplified Disease Activity Index.

† Drug retention rates were compared across the 3 groups with Kaplan-Meier with log rank test, continuous measures by Kruskal-Wallis test, and proportions by chi-square test or Fisher's exact test, as appropriate. Multiple comparisons between groups were conducted by log rank test, Kruskal-Wallis with post hoc Dunn test, chi-square test, or Fisher's exact test, as appropriate, with *P* values to be adjusted by applying the Holm's correction.

‡ Patients with at least 12 months from secukinumab start to date of data extraction. Patients who stopped treatment due to remission or other reasons (e.g., pregnancy) were censored at the stop date to reflect that their withdrawal was not due to lack of effectiveness or adverse events.

§ Statistically significant difference between b/tsDMARD-naïve patients and patients treated with ≥2 prior b/tsDMARDs.

¶ Statistically significant difference between b/tsDMARD-naïve patients and patients treated with 1 prior b/tsDMARD. Statistically significant difference between b/tsDMARD-naïve patients and patients treated with ≥2 prior b/tsDMARDs.

Statistically significant difference between b/tsDMARD-naïve patients and patients treated with 1 prior b/tsDMARD. Statistically significant difference between b/tsDMARD-naïve patients and patients treated with ≥2 prior b/tsDMARDs. Statistically significant difference between patients treated with 1 prior b/tsDMARD and ≥2 prior b/tsDMARDs. Significance level for all tests is 0.05.

(see Supplementary Tables 2 and 8). The number of included patients varied considerably across the European registries (from 30 to 657 patients). Significant differences in retention rates across the registries were observed, with 6-month retention rates varying between 80% (DANBIO) and 97% (TURKBIO), and 12-month retention rates varying from 51% (ROB-FIN) to 92% (RRBR and ATTRA) (Table 3 and Figure 2). Similar differences were found in adjusted analyses (see Supplementary Table 8).

Remission. Crude and LUNDEX-adjusted proportions of patients achieving DAPSA28, DAS28-CRP, SDAI, and CDAI remission after 6 and 12 months are presented in Table 2. DAPSA28, SDAI, and CDAI remission rates were similar (~10–15%), whereas approximately one-third of the patients achieved DAS28-CRP remission.

The proportion of patients achieving remission was significantly higher in biologics-naïve patients than in patients previously treated with 1 and ≥2 b/tsDMARDs (Table 2, Figure 3, and Supplementary Figure 1, available on the *Arthritis Care & Research* website at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560>). Adjusted analyses gave similar results (see Supplementary Table 9, available at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560>).

Crude and adjusted remission rates at 6 and 12 months of treatment were independent of time since diagnosis (see Supplementary Tables 2 and 9). Overall, heterogeneity in crude and adjusted remission rates across the European registries was found (Table 3 and Supplementary Table 7).

LDA (including remission). Crude and LUNDEX-adjusted proportions of patients achieving DAPSA28, DAS28-CRP, SDAI, and CDAI LDA after 6 and 12 months of

treatment are presented in Table 2, Figure 3, and Supplementary Figure 1, available at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560/abstract>. Overall, crude and LUNDEX-adjusted LDA rates were significantly higher in biologics-naïve patients, also in adjusted analyses (see Supplementary Table 9).

For all outcomes, achievement of LDA was independent of time since diagnosis (see Supplementary Table 2), also after adjustment (see Supplementary Table 9). Significant heterogeneities in crude (Table 3) and adjusted (see Supplementary Table 9) LDA rates were seen between the registries.

Response rates. ACR20/50/70 responses were achieved by 34%/19%/11% of the patients, and EULAR moderate/good response by 59% of the patients after 6 months. After 12 months, numbers were largely the same (Table 2). Changes in outcome measures from baseline to 6 months (and 12 months, respectively) were as follows: DAPSA28 –9.5 (–10.3), DAS28-CRP –0.9 (–1.1), SDAI –8.9 (–9.7), and CDAI –8.0 (–8.8).

Significantly better outcomes for ACR20/50/70 and EULAR moderate/good responses were observed for biologics-naïve patients (Table 2, Figure 3, and Supplementary Figure 1, available at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560/abstract>), also after adjustment (see Supplementary Table 9).

Response rates were independent of time since diagnosis (see Supplementary Table 2), also in adjusted analyses (see Supplementary Table 9). Significant heterogeneity in response rates between the European registries was found in crude as well as adjusted analyses (Table 3 and Supplementary Table 9).

Safety. Of the 2,017 patients starting secukinumab, 1,543 patients started treatment at least 12 months before date of data




Figure 1. **A**, Pooled 12-month secukinumab retention rates stratified by number of previous biologic/targeted synthetic disease-modifying antirheumatic drugs (b/tsDMARDs) (Kaplan-Meier curve with log rank test; $P = 0.001$). **B**, Cumulative incidence curve for withdrawal of secukinumab due to adverse events and lack of effectiveness.

extraction. Of these 1,543 patients, 602 patients withdrew from secukinumab before 12 months, of whom 107 patients withdrew due to adverse events. Time in weeks to secukinumab withdrawal for these 107 patients was similar across number of previous b/tsDMARDs (0/1/≥2) (Table 2). More patients withdrew from secukinumab due to lack of effectiveness than due to adverse events (Table 2). The cumulative incidence curve, which estimates the cumulative probabilities of treatment withdrawal over time, shows that the cumulative probability of withdrawal due to lack of effectiveness is higher than adverse events after ~4 months of treatment (Figure 1B).

Sensitivity analyses. Sensitivity analyses of 976 patients with ≥1 swollen joint (of 28) at the start of secukinumab treatment showed largely similar results to the analyses in Table 2 (see Supplementary Table 3, available at <http://onlinelibrary.wiley.com/doi/10.1002/acr.24560>). Sensitivity analyses of patients with secukinumab initiation at least 12 months before date of data extraction also showed largely similar results but did not reach significance for the 6-month comparison of retention rates across number of previous b/tsDMARDs (b/tsDMARD naïve: 89% [86–93%]; 1 prior

b/tsDMARD: 85% [81–89%]; ≥2 prior b/tsDMARDs: 85% [82–87%]; $P = 0.107$ [see Supplementary Table 4]).

DISCUSSION

This large real-life study of secukinumab effectiveness (i.e., drug retention, remission, LDA, and response rates) included 2,017 patients with PsA treated as part of routine care in 13 countries across Europe. Overall, high 6-month (86%) and 12-month (76%) secukinumab retention rates were found. Secukinumab effectiveness was significantly better for biologics-naïve patients after 6 as well as 12 months of treatment, was independent of time since diagnosis, and differed significantly across the European countries. Remission, LDA, and response rates were overall comparable to previous real-life observations in patients treated with a TNFi (5). Hence, this large observational study documents the effectiveness of secukinumab in the treatment of PsA patients.

Secukinumab effectiveness has previously been reported in one observational study of 76 Spanish PsA patients, in which 12-month retention rates were somewhat higher than in our study; for biologics-naïve patients, it was 91%, and for non-naïve patients, it was 82% (22). Good 1-year secukinumab effectiveness has also been reported in an Italian observational study of 130 PsA patients (23). In the FUTURE 1 RCT, 89% of the patients in the 150-mg secukinumab group reached 52 weeks, and ACR20/50 responses at week 24 and 52 were achieved by 50%/35% and 60%/43% of the patients, respectively (24). In our observational study, ACR20/50 responses at week 26 and 52 were lower than in the FUTURE 1 study (34%/19% and 37%/21%), probably reflecting that the study designs differed substantially (longitudinal observational study with 22% biologics-naïve patients versus RCT with 71% biologics-naïve patients). In the FUTURE 5 RCT, 91% of the patients treated with 150 mg of secukinumab completed 52 weeks of treatment, with ACR20/50/70 responses of 64%/41%/26%, thus substantially higher than in our study (10).

Interestingly, the overall secukinumab retention rates in this real-life study were similar to the retention rates of TNFi in a recently published observational study of 14,261 European biologics-naïve PsA patients (86% versus 86% at 6 months; 76% versus 77% at 12 months, respectively) and numerically slightly higher for biologics-naïve secukinumab than TNFi starters (90% versus 86% at 6 months, and 82% versus 77% at 12 months, respectively) (5). Overall, remission and response rates for patients treated with secukinumab were fairly similar to what was reported for TNFi (5) as well as to the effectiveness of TNFi reported in other, smaller observational studies (25–28).

Similar to findings in observational studies on TNFi, and in the FUTURE 2 and 5 trials, the current study demonstrated that effectiveness of secukinumab declines with increasing previous use of

Table 3. Retention, remission, low disease activity (including remission), and response rates after 6 and 12 months of secukinumab treatment across European observational registries (unadjusted analyses)*

	ARTIS (n = 657)	ATTRA (n = 151)	BADASER (n = 154)	BIO- BADASER (n = 154)	Biorx.si (n = 79)	DANBIO (n = 313)	GISEA (n = 180)	ICEBIO (n = 38)	NOR- DMARD (n = 60)	Reuma.pt (n = 68)	ROB-FIN (n = 47)	RRBR (n = 37)	SCQM (n = 203)	TURKBIO (n = 30)	P†
Drug retention rate, % (95% CI)															
6 months	82 (79-85)	94 (90-98)	93 (89-97)	92 (87-98)	80 (75-84)	96 (93-99)	87 (77-98)	83 (74-94)	91 (84-98)	83 (73-94)	92 (83-100)	90 (85-94)	97 (90-100)		<0.001
12 months	66 (62-70)	92 (88-97)	84 (78-91)	89 (82-96)	70 (65-76)	88 (82-93)	77 (64-93)	72 (61-86)	86 (78-96)	51 (39-68)	92 (83-100)	82 (77-88)	-		<0.001
DAPSA28 score ≤4															
6 months	8	22	-	11	12	-	0	14	14	19	-	35	21		<0.001
Crude	6	21	-	10	9	-	0	12	13	16	-	31	-		<0.001
LUNDEX															
12 months	6	23	-	10	14	-	NC	16	16	NC	-	15	NC		0.004
Crude	3	20	-	9	9	-	NC	11	14	NC	-	12	-		0.002
LUNDEX															
DAPSA28 score ≤14															
6 months	37	61	-	53	44	-	42	61	54	54	-	58	74		<0.001
Crude	30	58	-	49	33	-	35	50	49	45	-	51	-		<0.001
LUNDEX															
12 months	35	79	-	48	46	-	NC	68	63	NC	-	59	NC		<0.001
Crude	19	67	-	42	29	-	NC	49	55	NC	-	47	-		<0.001
LUNDEX															
DAS28-CRP score <2.6															
6 months	27	46	50	40	33	-	29	54	45	42	60	49	63		<0.001
Crude	21	44	44	37	25	-	24	44	41	35	52	43	-		<0.001
LUNDEX															
12 months	25	62	49	46	41	-	50	63	50	NC	NC	41	NC		<0.001
Crude	14	53	36	41	26	-	37	45	44	NC	NC	33	-		<0.001
LUNDEX															
DAS28-CRP score ≤3.2															
6 months	45	62	69	64	53	-	64	79	64	73	73	60	74		<0.001
Crude	36	58	60	59	40	-	54	64	58	61	64	53	-		<0.001
LUNDEX															
12 months	43	80	73	60	54	-	75	68	80	NC	NC	76	NC		<0.001
Crude	23	69	54	52	34	-	55	49	70	NC	NC	61	-		<0.001
LUNDEX															
SDAI score ≤3.3															
6 months	6	21	-	16	12	-	0	10	18	21	24	21	21		0.003
Crude	5	20	-	15	9	-	0	8	16	17	21	18	-		0.003
LUNDEX															
12 months	8	32	-	23	14	-	NC	25	5	NC	NC	15	NC		0.002
Crude	4	27	-	20	8	-	NC	18	4	NC	NC	12	-		<0.001
LUNDEX															

(Continued)

Table 3. (Cont'd)

	ARTIS (n = 657)	ATTRA (n = 151)	BIO- BADASER (n = 154)	Biorx.si (n = 79)	DANBIO (n = 313)	GISEA (n = 180)	ICEBIO (n = 38)	NOR- DMARD (n = 60)	Reuma.pt (n = 68)	ROB-FIN (n = 47)	RRBR (n = 37)	SCQM (n = 203)	TURKBIO (n = 30)	P†
SDAI score ≤11														
6 months														
Crude	42	68	-	58	54	-	46	67	64	67	76	65	74	<0.001
LUNDEX	33	64	-	53	41	-	39	54	58	55	66	57	-	<0.001
12 months														
Crude	50	88	-	56	55	-	NC	83	85	NC	NC	67	NC	<0.001
LUNDEX	27	75	-	49	35	-	NC	59	74	NC	NC	53	-	<0.001
CDAI score ≤2.8														
6 months														
Crude	6	18	-	12	14	-	0	9	11	12	20	23	21	0.007
LUNDEX	5	17	-	11	10	-	0	7	10	10	17	21	-	0.008
12 months														
Crude	8	31	-	19	14	-	14	25	10	NC	NC	23	NC	0.003
LUNDEX	4	27	-	17	9	-	10	18	9	NC	NC	18	-	<0.001
CDAI score ≤10														
6 months														
Crude	41	68	-	60	53	-	41	59	64	62	76	64	74	<0.001
LUNDEX	33	64	-	55	40	-	35	48	58	52	66	56	-	<0.001
12 months														
Crude	44	88	-	58	56	-	57	83	80	NC	NC	63	NC	<0.001
LUNDEX	24	75	-	51	35	-	42	59	70	NC	NC	50	-	<0.001
ACR/EULAR Boolean remission														
6 months														
Crude	5	22	18	9	9	0	0	8	6	12	23	15	16	<0.001
LUNDEX	4	21	15	8	7	0	0	7	6	10	20	13	-	<0.001
12 months														
Crude	5	24	15	7	9	0	5	10	8	15	NC	7	NC	<0.001
LUNDEX	3	21	11	7	6	0	4	7	7	8	NC	5	-	<0.001
ACR20 response														
6 months														
Crude	24	55	-	59	25	-	NC	NC	56	-	-	NC	22	<0.001
LUNDEX	20	51	-	54	19	-	NC	NC	51	-	-	18	-	<0.001
12 months														
Crude	27	67	-	50	24	-	NC	NC	NC	-	-	NC	NC	<0.001
LUNDEX	14	58	-	44	15	-	NC	NC	NC	-	-	24	-	<0.001
ACR50 response														
6 months														
Crude	11	36	-	38	12	-	NC	NC	31	-	-	NC	11	<0.001
LUNDEX	9	34	-	35	9	-	NC	NC	28	-	-	NC	-	<0.001
12 months														
Crude	15	45	-	35	11	-	NC	NC	NC	-	-	NC	NC	<0.001
LUNDEX	8	39	-	30	7	-	NC	NC	NC	-	-	NC	-	<0.001

(Continued)

Table 3. (Cont'd)

	ARTIS (n = 657)	ATTRA (n = 151)	BIO- BADASER (n = 154)	Biorx.si (n = 79)	DANBIO (n = 313)	GISEA (n = 180)	ICEBIO (n = 38)	NOR- DMARD (n = 60)	Reuma.pt (n = 68)	ROB-FIN (n = 47)	RRBR (n = 37)	SCQM (n = 203)	TURKBIO (n = 30)	P†
ACR70 response														
6 months														
Crude	6	21	-	21	7	-	NC	NC	19	-	-	NC	11	0.010
LUNDEX	4	20	-	19	6	-	NC	NC	17	-	-	NC	-	0.001
12 months														
Crude	5	24	-	31	5	-	NC	NC	NC	-	-	NC	NC	0.002
LUNDEX	3	20	-	27	3	-	NC	NC	NC	-	-	NC	-	0.001
EULAR good/moderate response														
6 months														
Crude	50	88	69	83	50	-	NC	55	62	47	93	NC	39	<0.001
LUNDEX	40	82	61	76	38	-	NC	45	56	39	81	NC	-	<0.001
12 months														
Crude	48	93	63	77	60	-	NC	43	83	NC	NC	64	NC	<0.001
LUNDEX	26	79	47	68	37	-	NC	30	73	NC	NC	50	-	<0.001

* Values are the percentage unless indicated otherwise. Numbers available for each of the analyses are shown in Supplementary Table 7. Registries and countries are as follows: ARTIS (Sweden), DANBIO (Denmark), SCQM (Switzerland), GISEA (Italy), BIOBADASER (Spain), ATTRA (Czech Republic), biorx.si (Slovenia), Reuma.pt (Portugal), NOR-DMARD (Norway), ROB-FIN (Finland), ICEBIO (Iceland), RRBR (Romania), and TURKBIO (Turkey). 95% CI = 95% confidence interval; ACR = American College of Rheumatology; ACR20/50/70 = ACR 20%/50%/70% improvement criteria; CDAl = Clinical Disease Activity Index; DAPSA28 = 28-joint Disease Activity Index for Psoriatic Arthritis; DAS28-CRP = Disease Activity Score in 28 joints using the C-reactive protein level; EULAR = European Alliance of Associations for Rheumatology; NC = not calculated (because data from <10 patients available); SDAI = Simplified Disease Activity Index.

† Comparisons between the registries were performed with Kaplan-Meier with log rank test for retention rates and chi-square test or Fisher's exact test for remission and response rates, as appropriate.




Figure 2. Twelve-month secukinumab retention rates compared across the European registries (Kaplan-Meier curve with log rank test; $P < 0.001$). Registries and countries are as follows: ARTIS (Sweden), DANBIO (Denmark), SCQM (Switzerland), GISEA (Italy), BIOBADASER (Spain), ATTRA (Czech Republic), biorx.si (Slovenia), Reuma.pt (Portugal), NOR-DMARD (Norway), ROB-FIN (Finland), ICEBIO (Iceland), RRBR (Romania), and TURKBIO (Turkey).

b/tsDMARDs, possibly reflecting confounding by indication (9,27,29,30). The similar secukinumab effectiveness for patients with different disease durations found in this study is also in accordance with previous findings for TNFi in patients with PsA (31–33).

In the 2017 updated treat-to-target recommendations for PsA, the DAPSA and minimal disease activity (MDA) are the preferred measures to define treatment target in PsA patients (34). In our study, the DAPSA (including a 66 swollen/68 tender joint count) (35) was only available in a minority of patients. Instead, we used the DAPSA28, which only requires a 28-joint count (13). The DAPSA28 has shown good criterion, correlational, and construct validity, as well as sensitivity to change, although the original DAPSA should be preferred when available (13). MDA was not assessed in the study due to lack of data on enthesitis and psoriasis in the majority of registries.

We chose the DAS28-CRP over the DAS28-ESR due to less missing data for the DAS28-CRP. Overall, the DAS28-CRP was a more liberal remission criterion than the SDAI, the CDAI, and the DAPSA28 in our study, which is consistent with previous reports (5,12,36,37). In the DAPSA28, SDAI, and CDAI LDA measures, we chose to include remission in accordance with the DAS28 LDA, as we believe that rheumatologists will be mainly interested in knowing how many patients at least were in LDA (i.e., in LDA or remission).

The major strength of this study is the 12-month longitudinal, observational study design with inclusion of a high number of PsA patients from 13 different countries. Furthermore, the data

included in the study were collected independently of commercial interests as part of standard care. Hence, although Novartis supports the EuroSpA collaboration, Novartis had no influence on data collection, statistical analyses, manuscript preparation, or the decision to submit. Major limitations of the study include lack of data on extraarticular inflammatory involvement and the fact that data on the optimal number of joints (66/68) were generally not available, which may have led to underestimation of disease activity. Furthermore, the DAS28, the CDAI, and the SDAI are composite scores originally developed for RA and not PsA.

Heterogeneity in baseline characteristics and secukinumab effectiveness across the registries was found. Importantly, the number of included patients (from 30 to 657) and proportions of biologics-naïve patients (from 5% to 97%) varied considerably across the registries and may explain some of the heterogeneity in effectiveness measures, e.g., a higher proportion of biologics-naïve patients may positively impact upon treatment outcomes. Moreover, low patient numbers in some registries will lead to more uncertain estimates, i.e., single patients will have a higher influence on outcomes. Also, the influence of different treatment guidelines and access to treatment in the different European countries were not accounted for in this study. Hence, interpretation of the pooled analyses should be done with caution. Of note, however, consistent results in prespecified unadjusted and adjusted analyses were found.

Furthermore, as is often the case in observational studies, some missing data on disease states and response rates were observed, challenging the generalizability of the findings.




Figure 3. Bar charts of crude proportions of patients achieving remission (A), LDA (including remission) (B), and response rates (C) after 12 months of secukinumab treatment compared across number of previous biologic/targeted synthetic disease-modifying antirheumatic drugs (b/tsDMARDs) for b/tsDMARDs naive (red), 1 prior b/tsDMARD (green), and ≥2 prior b/tsDMARDs (blue). ACR = American College of Rheumatology; CDAI = Clinical Disease Activity Index; DAPSA28 = 28-joint Disease Activity Index for Psoriatic Arthritis; DAS28-CRP = Disease Activity Score in 28 joints using the CRP level; EULAR = European Alliance of Associations for Rheumatology; SDAI = Simplified Disease Activity Index.

However, the study is by far the largest real-life study to date on secukinumab effectiveness in patients with PsA.

In conclusion, in this longitudinal observational study of >2,000 patients with PsA treated with secukinumab, we found high retention rates after 6 and 12 months of treatment and good remission, LDA, and response rates. Secukinumab effectiveness was significantly better for biologics-naïve patients, was independent of time since diagnosis, and varied across European registries.

AUTHOR CONTRIBUTIONS

All authors were involved in drafting the article or revising it critically for important intellectual content, and all authors approved the final version to be submitted for publication. Dr. Michelsen had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study conception and design. Michelsen, Georgiadis, Di Giuseppe, Loft, Nissen, Iannone, Pombo-Suarez, Mann, Rotar, Eklund, Kvien, Santos, Gudbjornsson, Codreanu, Yilmaz, Wallman, Brahe, Möller, Favalli, Sánchez-Piedra, Nekvindova, Tomsic, Trokovic, Kristianslund, Santos, Löve, Ionescu, Pehlivan, Jones, van der Horst-Bruinsma, Ørnberg, Østergaard, Hetland.

Acquisition of data. Michelsen, Georgiadis, Di Giuseppe, Loft, Nissen, Iannone, Pombo-Suarez, Mann, Rotar, Eklund, Kvien, Santos, Gudbjornsson, Codreanu, Yilmaz, Wallman, Brahe, Möller, Favalli, Sánchez-Piedra, Nekvindova, Tomsic, Trokovic, Kristianslund, Santos, Löve, Ionescu, Pehlivan, Jones, van der Horst-Bruinsma, Ørnberg, Østergaard, Hetland.

Analysis and interpretation of data. Michelsen, Georgiadis, Di Giuseppe, Loft, Nissen, Iannone, Pombo-Suarez, Mann, Rotar, Eklund, Kvien, Santos, Gudbjornsson, Codreanu, Yilmaz, Wallman, Brahe, Möller, Favalli, Sánchez-Piedra, Nekvindova, Tomsic, Trokovic, Kristianslund, Santos, Löve, Ionescu, Pehlivan, Jones, van der Horst-Bruinsma, Ørnberg, Østergaard, Hetland.

ROLE OF THE STUDY SPONSOR

Novartis had no role in the study design or in the collection, analysis, or interpretation of the data, the writing of the manuscript, or the decision to submit the manuscript for publication. Publication of this article was not contingent upon approval by Novartis.

REFERENCES

- Gladman DD, Antoni C, Mease P, Clegg DO, Nash P. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. *Ann Rheum Dis* 2005;64 Suppl 2:ii14–7.
- Michelsen B, Diamantopoulos AP, Hoiberg HK, Soldal DM, Kavanaugh A, Haugeberg G. Need for improvement in current treatment of psoriatic arthritis: study of an outpatient clinic population. *J Rheumatol* 2017;44:431–6.
- Michelsen B, Uhlig T, Sexton J, van der Heijde D, Hammer HB, Kristianslund EK, et al. Health-related quality of life in patients with psoriatic and rheumatoid arthritis: data from the prospective multicentre nor-DMARD study compared with Norwegian general population controls. *Ann Rheum Dis* 2018;77:1290–4.
- Gossec L, Baraliakos X, Kerschbaumer A, de Wit M, McInnes I, Dougados M, et al. EULAR recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. *Ann Rheum Dis* 2020;79:700–12.
- Brahe CH, Ørnberg LM, Jacobsson L, Nissen MJ, Kristianslund EK, Mann H, et al. Retention and response rates in 14 261 PsA patients starting TNF inhibitor treatment—results from 12 countries in EuroSpA. *Rheumatology (Oxford)* 2020;59:1640–50.
- Hetland ML. Psoriatic arthritis: still room for improvement. *Lancet* 2020;395:1463–5.
- McInnes IB, Mease PJ, Kirkham B, Kavanaugh A, Ritchlin CT, Rahman P, et al. Secukinumab, a human anti-interleukin-17a monoclonal antibody, in patients with psoriatic arthritis (future 2): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2015;386:1137–46.
- McInnes IB, Mease PJ, Ritchlin CT, Rahman P, Gottlieb AB, Kirkham B, et al. Secukinumab sustains improvement in signs and symptoms

- of psoriatic arthritis: 2 year results from the phase 3 future 2 study. *Rheumatology (Oxford)* 2017;56:1993–2003.
9. Mease P, van der Heijde D, Landewe R, Mpofu S, Rahman P, Tahir H, et al. Secukinumab improves active psoriatic arthritis symptoms and inhibits radiographic progression: primary results from the randomised, double-blind, phase III future 5 study. *Ann Rheum Dis* 2018;77:890–7.
 10. Van der Heijde D, Mease PJ, Landewe RB, Rahman P, Tahir H, Singhal A, et al. Secukinumab provides sustained low rates of radiographic progression in psoriatic arthritis: 52-week results from a phase 3 study, future 5. *Rheumatology (Oxford)* 2020;59:1325–34.
 11. Ornbjerg LM, Brahe CH, Askling J, Ciurea A, Mann H, Onen F, et al. Treatment response and drug retention rates in 24 195 biologic-naïve patients with axial spondyloarthritis initiating TNF treatment: routine care data from 12 registries in the EuroSpA collaboration. *Ann Rheum Dis* 2019;78:1536–44.
 12. Michelsen B, Ornbjerg LM, Kvien TK, Pavelka K, Nissen MJ, Nordstrom D, et al. Impact of discordance between patient's and evaluator's global assessment on treatment outcomes in 14 868 patients with spondyloarthritis. *Rheumatology (Oxford)* 2020;59:2455–61.
 13. Michelsen B, Sexton J, Smolen JS, Aletaha D, Krogh NS, van der Heijde D, et al. Can disease activity in patients with psoriatic arthritis be adequately assessed by a modified disease activity index for psoriatic arthritis (DAPSA) based on 28 joints? *Ann Rheum Dis* 2018;77:1736–41.
 14. Wells G, Becker JC, Teng J, Dougados M, Schiff M, Smolen J, et al. Validation of the 28-joint Disease Activity Score (DAS28) and European League Against Rheumatism response criteria based on C-reactive protein against disease progression in patients with rheumatoid arthritis, and comparison with the DAS28 based on erythrocyte sedimentation rate. *Ann Rheum Dis* 2009;68:954–60.
 15. Aletaha D, Smolen JS. The Simplified Disease Activity Index (SDAI) and Clinical Disease Activity Index (CDAI) to monitor patients in standard clinical care. *Best Pract Res Clin Rheumatol* 2007;21:663–75.
 16. Smolen JS, Braun J, Dougados M, Emery P, Fitzgerald O, Helliwell P, et al. Treating spondyloarthritis, including ankylosing spondylitis and psoriatic arthritis, to target: recommendations of an international task force. *Ann Rheum Dis* 2014;73:6–16.
 17. Fransen J, Antoni C, Mease PJ, Uter W, Kavanaugh A, Kalden JR, et al. Performance of response criteria for assessing peripheral arthritis in patients with psoriatic arthritis: analysis of data from randomised controlled trials of two tumour necrosis factor inhibitors. *Ann Rheum Dis* 2006;65:1373–8.
 18. Felson DT, Smolen JS, Wells G, Zhang B, van Tuyl LH, Funovits J, et al. American College of Rheumatology/European League Against Rheumatism provisional definition of remission in rheumatoid arthritis for clinical trials. *Arthritis Rheum* 2011;63:573–86.
 19. Felson DT, Anderson JJ, Boers M, Bombardier C, Furst D, Goldsmith C, et al. American College of Rheumatology preliminary definition of improvement in rheumatoid arthritis. *Arthritis Rheum* 1995;38:727–35.
 20. Kristensen LE, Saxne T, Geborek P. The LUNDEX, a new index of drug efficacy in clinical practice: results of a five-year observational study of treatment with infliximab and etanercept among rheumatoid arthritis patients in southern Sweden. *Arthritis Rheum* 2006;54:600–6.
 21. Dixon WG, Carmona L, Finckh A, Hetland ML, Kvien TK, Landewe R, et al. EULAR points to consider when establishing, analysing and reporting safety data of biologics registers in rheumatology. *Ann Rheum Dis* 2010;69:1596–602.
 22. Pinto Tasende JA, Maceiras Pan FJ, Mosquera Martinez JA, Fernandez Dominguez L, Correa Rey B, Garcia Porrua C. Secukinumab as biological treatment for psoriatic arthritis in real clinical practice. *Reumatol Clin (Engl Ed)* 2021;17:203–6.
 23. Chimenti MS, Fonti GL, Conigliaro P, Sunzini F, Scrivo R, Navarini L, et al. One-year effectiveness, retention rate, and safety of secukinumab in ankylosing spondylitis and psoriatic arthritis: a real-life multicenter study. *Expert Opin Biol Ther* 2020;20:813–21.
 24. Mease PJ, McInnes IB, Kirkham B, Kavanaugh A, Rahman P, van der Heijde D, et al. Secukinumab inhibition of interleukin-17a in patients with psoriatic arthritis. *N Engl J Med* 2015;373:1329–39.
 25. Glinthorg B, Østergaard M, Dreyer L, Krogh NS, Tarp U, Hansen MS, et al. Treatment response, drug survival, and predictors thereof in 764 patients with psoriatic arthritis treated with anti-tumor necrosis factor α therapy: results from the nationwide Danish DANBIO registry. *Arthritis Rheum* 2011;63:382–90.
 26. Stober C, Ye W, Guruparan T, Htut E, Clunie G, Jadon D. Prevalence and predictors of tumour necrosis factor inhibitor persistence in psoriatic arthritis. *Rheumatology (Oxford)* 2018;57:158–63.
 27. Fagerli KM, Lie E, van der Heijde D, Heiberg MS, Kalstad S, Rodevand E, et al. Switching between TNF inhibitors in psoriatic arthritis: data from the nor-DMARD study. *Ann Rheum Dis* 2013;72:1840–4.
 28. Carmona L, Gomez-Reino JJ. Survival of TNF antagonists in spondyloarthritis is better than in rheumatoid arthritis: data from the Spanish registry BIOBADASER. *Arthritis Res Ther* 2006;8:R72.
 29. Costa L, Perricone C, Chimenti MS, Del Puente A, Caso P, Peluso R, et al. Switching between biological treatments in psoriatic arthritis: a review of the evidence. *Drugs R D* 2017;17:509–22.
 30. Kavanaugh A, McInnes IB, Mease PJ, Hall S, Chinoy H, Kivitz AJ, et al. Efficacy of subcutaneous secukinumab in patients with active psoriatic arthritis stratified by prior tumor necrosis factor inhibitor use: results from the randomized placebo-controlled future 2 study. *J Rheumatol* 2016;43:1713–7.
 31. Carvalho PD, Duarte C, Vieira-Sousa E, Cunha-Miranda L, Avila-Ribeiro P, Santos H, et al. Predictors of response to TNF blockers in patients with polyarticular psoriatic arthritis. *Acta Reumatol Port* 2017;42:55–65.
 32. Fagerli KM, Lie E, van der Heijde D, Heiberg MS, Lexberg AS, Rodevand E, et al. The role of methotrexate co-medication in TNF-inhibitor treatment in patients with psoriatic arthritis: results from 440 patients included in the nor-DMARD study. *Ann Rheum Dis* 2014;73:132–7.
 33. Perrotta FM, Marchesoni A, Lubrano E. Minimal disease activity and remission in psoriatic arthritis patients treated with anti-TNF- α drugs. *J Rheumatol* 2016;43:350–5.
 34. Smolen JS, Schols M, Braun J, Dougados M, Fitzgerald O, Gladman DD, et al. Treating axial spondyloarthritis and peripheral spondyloarthritis, especially psoriatic arthritis, to target: 2017 update of recommendations by an international task force. *Ann Rheum Dis* 2018;77:3–17.
 35. Schoels M, Aletaha D, Funovits J, Kavanaugh A, Baker D, Smolen JS. Application of the DAREA/DAPSA score for assessment of disease activity in psoriatic arthritis. *Ann Rheum Dis* 2010;69:1441–7.
 36. Fleischmann R, van der Heijde D, Koenig AS, Pedersen R, Szumski A, Marshall L, et al. How much does disease activity score in 28 joints ESR and CRP calculations underestimate disease activity compared with the simplified disease activity index? *Ann Rheum Dis* 2015;74:1132–7.
 37. Michelsen B, Kristianslund EK, Sexton J, Hammer HB, Fagerli KM, Lie E, et al. Do depression and anxiety reduce the likelihood of remission in rheumatoid arthritis and psoriatic arthritis? Data from the prospective multicentre nor-DMARD study. *Ann Rheum Dis* 2017;76:1906–10.